



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\frac{1}{2} \begin{vmatrix} a_2, & b_2, & a_2, & b_2 \\ a_2, & b_3, & a_3, & b_3 \\ a_4, & b_4, & a_4, & b_4 \\ a_5, & b_5, & a_5, & b_5 \end{vmatrix} \equiv (23)(45) - (24)(35) + (25)(34) \equiv 0,$$

which cannot be satisfied by

$$(23) = (45) = (24) = (35) = (25) = (34).$$

The general case fails in consequence of a similar identical relation among the determinants of a matrix. This relation can be expressed as the expansion of a determinant of $2(n-3)$ rows which can be symbolized by $\frac{A}{B} \bigg| \frac{C}{O}$, where A , B , C denote, respectively,

$$\begin{vmatrix} 0, & 0, & \dots, & a_{n-2} \\ 0, & 0, & \dots, & b_{n-2} \\ \dots & \dots & \dots & \dots \\ 0, & 0, & \dots, & l_{n-2} \end{vmatrix}, \quad \begin{vmatrix} a_2, & a_3, & a_4, & \dots, & a_{n-2} \\ b_2, & b_3, & b_4, & \dots, & b_{n-2} \\ \dots & \dots & \dots & \dots & \dots \\ l_2, & l_3, & l_4, & \dots, & l_{n-2} \end{vmatrix}, \quad \begin{vmatrix} a_{n-1}, & a_n, & a_2, & a_3, & \dots, & a_{n-4} \\ b_{n-1}, & b_n, & b_2, & b_3, & \dots, & b_{n-4} \\ \dots & \dots & \dots & \dots & \dots & \dots \\ l_{n-1}, & l_n, & l_2, & l_3, & \dots, & l_{n-4} \end{vmatrix}.$$

When this is expanded according to Laplace's method in determinants of $(n-3)$ rows, we get

$$\begin{aligned} & (2, 3, \dots, n-3, n-2)(n-1, n, 2, \dots, n-4) \\ & - (2, 3, \dots, n-3, n-1)(n-2, n, 2, \dots, n-4) \\ & + (2, 3, \dots, n-3, n)(n-3, n, 2, \dots, n-4) = 0, \end{aligned}$$

which cannot be satisfied if these determinants are all equal.

DEPARTMENTS.

SOLUTIONS OF PROBLEMS.

ALGEBRA.

Problem 206 was also solved by J. E. Sanders; No. 207 was also solved by L. E. Newcomb.

208. Proposed by F. P. MATZ, Sc. D., Ph. D., Professor of Mathematics and Astronomy in Defiance College, Defiance, O.

Solve (1)..... $x^4 + y^4 = 14x^2y^2$; (2)..... $x + y = m$.

I. Solution by EDWIN L. RICH, Student at Lehigh University.

Equation (1) may be written

$$(x^2 - y^2 - xy\sqrt{12})(x^2 - y^2 + xy\sqrt{12}) = 0 \dots\dots (3).$$

Solving each factor,

$$x = (\sqrt{3} \pm 2)y \dots\dots (4),$$

$$x = (-\sqrt{3} \pm 2)y \dots\dots (5).$$

From the simultaneous equations (2), (4) and (2), (5),

$$x = \frac{m}{6}(3 \pm \sqrt{3}), \quad y = \frac{m}{6}(3 \mp \sqrt{3}),$$

$$x = \frac{m}{2}(1 \pm \sqrt{3}), \quad y = \frac{m}{2}(1 \mp \sqrt{3}).$$

II. Solution by S. A. COREY, Hitsman, Iowa.

Adding $2x^2y^2$ to each member of (1),

$$x^4 + 2x^2y^2 + y^4 = 16x^2y^2 \dots\dots (3),$$

whence

$$x^2 + y^2 = \pm 4xy \dots\dots (4).$$

Squaring (2), we have

$$x^2 + 2xy + y^2 = m^2 \dots\dots (5);$$

substituting from (4),

$$6xy = m^2, \text{ or } -2xy = m^2,$$

whence,

$$x = \frac{m^2}{6y} \text{ or } -\frac{m^2}{2y}.$$

By substituting these values of x in (2), we find without difficulty,

$$x = m(\frac{1}{2} \pm \sqrt{\frac{1}{12}}), \text{ or } m(\frac{1}{2} \pm \sqrt{3}/4);$$

$$y = m(\frac{1}{2} \mp \sqrt{\frac{1}{12}}), \text{ or } m(\frac{1}{2} \mp \sqrt{3}/4).$$

Also solved by R. L. Borger, G. W. Greenwood, E. L. Sherwood, L. E. Newcomb, J. F. Lawrence, S. E. Harwood, Elmer Schuyler, G. B. M. Zerr, and J. Scheffer.

209. Proposed by F. P. MATZ, Sc. D., Ph. D., Professor of Mathematics and Astronomy in Defiance College, Defiance, O.

Prove that $(a^4 + b^4 + c^4 + d^4) > 4abcd$.

Solution by M. E. GRABER, A. M., Heidelberg University, Tiffin, O.

$(a^2 + b^2) > 2ab$ and $(c^2 + d^2) > 2cd$ from which, by multiplication, $(a^2c^2 + b^2c^2 + a^2d^2 + b^2d^2) > 4abcd \dots\dots (1)$.

Again $(a^4 + c^4) > 2a^2c^2$; $(b^4 + c^4) > 2b^2c^2$; $(a^4 + d^4) > 2a^2d^2$, and $(b^4 + d^4) > 2b^2d^2$.

By addition, $(a^4 + b^4 + c^4 + d^4) > (a^2c^2 + b^2c^2 + a^2d^2 + b^2d^2)$ and this in connection with (1) gives $(a^4 + b^4 + c^4 + d^4) > 4abcd$.

Also solved by G. W. Greenwood, R. L. Borger, E. L. Sherwood, L. E. Newcomb, Elmer Schuyler, J. H. Meyer, A. J. Haun, O. A. Laisant, J. Scheffer.